

## GCACH MTD Toolkit: P4P Reporting Metrics

1 • Antidepressant Medication Management* ⌘	2A			
2 • Child and Adolescents' Access to Primary Care Practitioners*	2A	3D		
3 • Comprehensive Diabetes Care: Eye Exam (retinal) performed	2A	3D		
4 • Comprehensive Diabetes Care: Hemoglobin A1c Testing	2A	3D		
5 • Comprehensive Diabetes Care: Medical Attention for Nephropathy	2A	3D		
6 • Follow-up After Discharge from ED for Mental Health, Alcohol or Other Drug Dependence*	2A	2C		
7 • Follow-up After Hospitalization for Mental Illness*	2A	2C		
8 • Inpatient Hospital Utilization	2A	2C	3A	3D
9 • Medication Management for People with Asthma (5 – 64 Years) ⌘	2A	3D		
10 • Mental Health Treatment Penetration (Broad Version)* ⌘	2A			
11 • Outpatient Emergency Department Visits per 1000 member months* ⌘	2A	2C	3A	3D
12 • Patients on high-dose chronic opioid therapy by varying thresholds*	3A			
13 • Patients with concurrent sedatives prescriptions	3A			
14 • Percent Homeless (Narrow definition)*	2C			
15 • Plan All-Cause Readmission Rate (30 Days) ⌘	2A	2C		
16 • Statin Therapy for Patients with Cardiovascular Disease (Prescribed)	3D			
17 • Substance Use Disorder Treatment Penetration* ⌘	2A		3A	

2A Bi-Directional Integration

2C Transitional Care

3A Opioid Use

3D Chronic Disease

\* Measure contains sub-measures ⌘ ACH High Performance Metrics (QIS)

Updated: 12/4/2017

## GCACH Pay-For-Performance (P4P) Measure Set

Measure Name	Measure Description	Statewide DSRIP Accountability Measure?	Method for Assessment of ACH Performance	Initial Assessment Year	Associated Project Areas
Antidepressant Medication Management	The percentage of Medicaid enrollees 18 years of age and older with a diagnosis of major depression and were newly treated with antidepressant medication, and who remained on an antidepressant medication treatment.	Yes	Gap to Goal	2019	2A. Bi-Directional Integration
Child and Adolescents' Access to Primary Care Practitioners	Percent of children enrolled in Medicaid who had a visit with a primary care provider. Reported separately for the following age groups: <ul style="list-style-type: none"> <li>• 12-24 months,</li> <li>• 2-6 years,</li> <li>• 7-11 years, and</li> <li>• 12-19 years</li> </ul>	No	Gap to Goal	2019	2A. Bi-Directional Integration 3D. Chronic Disease Prevention
Comprehensive Diabetes Care: Eye Exam (retinal) performed	Percentage of Medicaid enrollees 18-75 years of age with diabetes who had a retinal or dilated eye exam by an eye care professional during the measurement period or a negative retinal exam (no evidence of retinopathy) in the 12 months prior to the measurement period.	No	Gap to Goal	2020	2A. Bi-Directional Integration 3D. Chronic Disease Prevention
Comprehensive Diabetes Care: Hemoglobin A1c Testing	The percentage of Medicaid enrollees 18–75 years of age with diabetes (type 1 and type 2) who received an HbA1c test during the measurement year.	No	Gap to Goal	2019	2A. Bi-Directional Integration 3D. Chronic Disease Prevention

Measure Name	Measure Description	Statewide DSRIP Accountability Measure?	Method for Assessment of ACH Performance	Initial Assessment Year	Associated Project Areas
Comprehensive Diabetes Care: Medical Attention for Nephropathy	The percentage of Medicaid enrollees 18–75 years of age with diabetes who had a nephropathy screening test or evidence of nephropathy during the measurement period.	No	Gap to Goal	2019	2A. Bi-Directional Integration 3D. Chronic Disease Prevention
Follow-up After Discharge from ED for Mental Health	The percentage of emergency department (ED) visits for beneficiaries with a principal diagnosis of mental illness, who had a follow up visit for mental illness within 30 days of discharge. Two rates are reported: (1) The percentage of discharges for beneficiaries who received follow-up within 30 days of discharge; (2) The percentage of discharges for beneficiaries who received follow-up within 7 days of discharge.	No	Improvement Over Self	2020	2A. Bi-Directional Integration 2C. Transitional Care
Follow-up After Discharge from ED for Alcohol or Other Drug Dependence	The percentage of emergency department (ED) visits for beneficiaries with a principal diagnosis of alcohol or other drug (AOD) dependence, who had a follow up visit for AOD within 30 days of the ED visit. Two rates are reported: (1) The percentage of discharges for beneficiaries who received follow-up within 30 days of discharge; (2) The	No	Improvement Over Self	2020	2A. Bi-Directional Integration 2C. Transitional Care

Measure Name	Measure Description	Statewide DSRIP Accountability Measure?	Method for Assessment of ACH Performance	Initial Assessment Year	Associated Project Areas
	percentage of discharges for beneficiaries who received follow-up within 7 days of discharge.				
Follow-up After Hospitalization for Mental Illness	The percentage of discharges for Medicaid enrollees 6 years of age and older who were hospitalized for treatment of selected mental illness diagnoses and who had an outpatient visit, an intensive outpatient encounter or partial hospitalization with a mental health practitioner. Two rates are reported: (1) The percentage of discharges for enrollees who received follow-up within 30 days of discharge; (2) The percentage of discharges the enrollees who received follow-up within 7 days of discharge.	No	Improvement Over Self	2020	2A. Bi-Directional Integration 3C. Transitional Care
Inpatient Hospital Utilization	For members 18 years of age and older, the risk-adjusted ratio of observed to expected acute inpatient discharges during the measurement year reported by Surgery, Medicine and Total.	No	Improvement Over Self	2020	2A. Bi-Directional Integration 2C. Transitional Care 3A. Opioid Use 3D. Chronic Disease Prevention
Medication Management for People with Asthma (5 – 64 Years)	The percentage of Medicaid enrollees 5-64 years of age during the measurement year who were identified as having persistent asthma and were dispensed appropriate medications that they	Yes	Gap to Goal	2019	2A. Bi-Directional Integration 3D. Chronic Disease Prevention

Measure Name	Measure Description	Statewide DSRIP Accountability Measure?	Method for Assessment of ACH Performance	Initial Assessment Year	Associated Project Areas
	remained on during the treatment period.				
Mental Health Treatment Penetration (Broad Version)	Percent of Medicaid enrollees with a mental health service need who received at least one qualifying service during the measurement year. Separate reporting by age groups: 12-17 years; 18-64 years; 65+ years.	Yes	Improvement Over Self	2019	2A. Bi-Directional Integration
Outpatient Emergency Department Visits per 1000 Member Months	The rate of Medicaid enrollee visits to emergency department per 1000 member months, including visits related to mental health and chemical dependency. Separate reporting for age groups 10-17, 18-64, and 65+.	Yes	Improvement Over Self	2019	2A. Bi-Directional Integration 2C. Transitional Care 3A. Opioid Use 3D. Chronic Disease Prevention
Patients on high-dose chronic opioid therapy by varying thresholds	Measure specification in development. Among Medicaid enrollees, the percentage of chronic opioid therapy patients receiving doses: >50 mg. MED in a quarter, doses >90 mg. MED in a quarter.	No	Improvement Over Self	2019	3A. Opioid Use
Patients with concurrent sedatives prescriptions	Measure specification in development. Among Medicaid enrollees receiving chronic opioid therapy, the percentage that had more than 45 days of Sedative Hypnotics/ Benzodiazepines/	No	Improvement Over Self	2019	3A. Opioid Use

Measure Name	Measure Description	Statewide DSRIP Accountability Measure?	Method for Assessment of ACH Performance	Initial Assessment Year	Associated Project Areas
	carisoprodol/ barbiturates dispensed in the quarter.				
Percent Homeless (Narrow Definition)	Percent of Medicaid enrollees who were homeless in at least one month in the measurement year. Excludes “homeless with housing” ACES living arrangement code	No	Improvement Over Self	2019	2C. Transitional Care
Plan All-Cause Readmission Rate (30 Days)	The proportion of acute inpatient stays during the measurement year that were followed by an unplanned acute readmission within 30 days among Medicaid enrollees ages 18-64 years old.	Yes	Improvement Over Self	2019	2A. Bi-Directional Integration 2C. Transitional Care
Statin Therapy for Patients with Cardiovascular Disease (Prescribed)	Percentage of male Medicaid enrollees 21 to 75 years of age and female Medicaid enrollees 40 to 75 years of age during the measurement year who were identified as having clinical ASCVD who were dispensed at least one high- or moderate-intensity statin medication.	No	Improvement Over Self	2020	3D. Chronic Disease Prevention
Substance Use Disorder Treatment Penetration	The percentage of Medicaid enrollees with a substance use disorder treatment need who received substance use disorder treatment in the measurement year Separate reporting by age groups: <ul style="list-style-type: none"> <li>• 12-17 years;</li> <li>• 18-64 years;</li> </ul>	Yes	Improvement Over Self	2019	2A. Bi-Directional Integration

Measure Name	Measure Description	Statewide DSRIP Accountability Measure?	Method for Assessment of ACH Performance	Initial Assessment Year	Associated Project Areas
	<ul style="list-style-type: none"> <li>65+ years.</li> </ul>				
Substance Use Disorder Treatment Penetration (Opioid)	Measure specification in development. Percent of Medicaid enrollees with a diagnosis of opioid use disorder who have a substance use service need who received at least one qualifying service during the measurement year. Reported separately for adults and for children.	No	Improvement Over Self	2020	3A. Opioid Use