

GCACH MTD Toolkit: P4P Reporting Metrics

| | | | | |
|--|----|----|----|----|
| 1 • Antidepressant Medication Management* ⌘ | 2A | | | |
| 2 • Child and Adolescents' Access to Primary Care Practitioners* | 2A | 3D | | |
| 3 • Comprehensive Diabetes Care: Eye Exam (retinal) performed | 2A | 3D | | |
| 4 • Comprehensive Diabetes Care: Hemoglobin A1c Testing | 2A | 3D | | |
| 5 • Comprehensive Diabetes Care: Medical Attention for Nephropathy | 2A | 3D | | |
| 6 • Follow-up After Discharge from ED for Mental Health, Alcohol or Other Drug Dependence* | 2A | 2C | | |
| 7 • Follow-up After Hospitalization for Mental Illness* | 2A | 2C | | |
| 8 • Inpatient Hospital Utilization | 2A | 2C | 3A | 3D |
| 9 • Medication Management for People with Asthma (5 – 64 Years) ⌘ | 2A | 3D | | |
| 10 • Mental Health Treatment Penetration (Broad Version)* ⌘ | 2A | | | |
| 11 • Outpatient Emergency Department Visits per 1000 member months* ⌘ | 2A | 2C | 3A | 3D |
| 12 • Patients on high-dose chronic opioid therapy by varying thresholds* | 3A | | | |
| 13 • Patients with concurrent sedatives prescriptions | 3A | | | |
| 14 • Percent Homeless (Narrow definition)* | 2C | | | |
| 15 • Plan All-Cause Readmission Rate (30 Days) ⌘ | 2A | 2C | | |
| 16 • Statin Therapy for Patients with Cardiovascular Disease (Prescribed) | 3D | | | |
| 17 • Substance Use Disorder Treatment Penetration* ⌘ | 2A | 3A | | |

2A Bi-Directional Integration

2C Transitional Care

3A Opioid Use

3D Chronic Disease

* Measure contains sub-measures ⌘ ACH High Performance Metrics (QIS)

Updated: 12/4/2017

GCACH Pay-For-Performance (P4P) Measure Set

| Measure Name | Measure Description | Statewide DSRIP Accountability Measure? | Method for Assessment of ACH Performance | Initial Assessment Year | Associated Project Areas |
|---|--|---|--|-------------------------|--|
| Antidepressant Medication Management | The percentage of Medicaid enrollees 18 years of age and older with a diagnosis of major depression and were newly treated with antidepressant medication, and who remained on an antidepressant medication treatment. | Yes | Gap to Goal | 2019 | 2A. Bi-Directional Integration |
| Child and Adolescents' Access to Primary Care Practitioners | Percent of children enrolled in Medicaid who had a visit with a primary care provider. Reported separately for the following age groups: <ul style="list-style-type: none"> • 12-24 months, • 2-6 years, • 7-11 years, and • 12-19 years | No | Gap to Goal | 2019 | 2A. Bi-Directional Integration 3D. Chronic Disease Prevention |
| Comprehensive Diabetes Care: Eye Exam (retinal) performed | Percentage of Medicaid enrollees 18-75 years of age with diabetes who had a retinal or dilated eye exam by an eye care professional during the measurement period or a negative retinal exam (no evidence of retinopathy) in the 12 months prior to the measurement period. | No | Gap to Goal | 2020 | 2A. Bi-Directional Integration 3D. Chronic Disease Prevention |
| Comprehensive Diabetes Care: Hemoglobin A1c Testing | The percentage of Medicaid enrollees 18–75 years of age with diabetes (type 1 and type 2) who received an HbA1c test during the measurement year. | No | Gap to Goal | 2019 | 2A. Bi-Directional Integration 3D. Chronic Disease Prevention |

| Measure Name | Measure Description | Statewide DSRIP Accountability Measure? | Method for Assessment of ACH Performance | Initial Assessment Year | Associated Project Areas |
|--|---|---|--|-------------------------|--|
| Comprehensive Diabetes Care: Medical Attention for Nephropathy | The percentage of Medicaid enrollees 18–75 years of age with diabetes who had a nephropathy screening test or evidence of nephropathy during the measurement period. | No | Gap to Goal | 2019 | 2A. Bi-Directional Integration 3D. Chronic Disease Prevention |
| Follow-up After Discharge from ED for Mental Health | The percentage of emergency department (ED) visits for beneficiaries with a principal diagnosis of mental illness, who had a follow up visit for mental illness within 30 days of discharge. Two rates are reported: (1) The percentage of discharges for beneficiaries who received follow-up within 30 days of discharge; (2) The percentage of discharges for beneficiaries who received follow-up within 7 days of discharge. | No | Improvement Over Self | 2020 | 2A. Bi-Directional Integration 2C. Transitional Care |
| Follow-up After Discharge from ED for Alcohol or Other Drug Dependence | The percentage of emergency department (ED) visits for beneficiaries with a principal diagnosis of alcohol or other drug (AOD) dependence, who had a follow up visit for AOD within 30 days of the ED visit. Two rates are reported: (1) The percentage of discharges for beneficiaries who received follow-up within 30 days of discharge; (2) The | No | Improvement Over Self | 2020 | 2A. Bi-Directional Integration 2C. Transitional Care |

| Measure Name | Measure Description | Statewide DSRIP Accountability Measure? | Method for Assessment of ACH Performance | Initial Assessment Year | Associated Project Areas |
|---|---|---|--|-------------------------|---|
| | percentage of discharges for beneficiaries who received follow-up within 7 days of discharge. | | | | |
| Follow-up After Hospitalization for Mental Illness | The percentage of discharges for Medicaid enrollees 6 years of age and older who were hospitalized for treatment of selected mental illness diagnoses and who had an outpatient visit, an intensive outpatient encounter or partial hospitalization with a mental health practitioner. Two rates are reported: (1) The percentage of discharges for enrollees who received follow-up within 30 days of discharge; (2) The percentage of discharges the enrollees who received follow-up within 7 days of discharge. | No | Improvement Over Self | 2020 | 2A. Bi-Directional Integration 3C. Transitional Care |
| Inpatient Hospital Utilization | For members 18 years of age and older, the risk-adjusted ratio of observed to expected acute inpatient discharges during the measurement year reported by Surgery, Medicine and Total. | No | Improvement Over Self | 2020 | 2A. Bi-Directional Integration 2C. Transitional Care 3A. Opioid Use 3D. Chronic Disease Prevention |
| Medication Management for People with Asthma (5 – 64 Years) | The percentage of Medicaid enrollees 5-64 years of age during the measurement year who were identified as having persistent asthma and were dispensed appropriate medications that they | Yes | Gap to Goal | 2019 | 2A. Bi-Directional Integration 3D. Chronic Disease Prevention |

| Measure Name | Measure Description | Statewide DSRIP Accountability Measure? | Method for Assessment of ACH Performance | Initial Assessment Year | Associated Project Areas |
|--|--|---|--|-------------------------|---|
| | remained on during the treatment period. | | | | |
| Mental Health Treatment Penetration (Broad Version) | Percent of Medicaid enrollees with a mental health service need who received at least one qualifying service during the measurement year. Separate reporting by age groups: 12-17 years; 18-64 years; 65+ years. | Yes | Improvement Over Self | 2019 | 2A. Bi-Directional Integration |
| Outpatient Emergency Department Visits per 1000 Member Months | The rate of Medicaid enrollee visits to emergency department per 1000 member months, including visits related to mental health and chemical dependency. Separate reporting for age groups 10-17, 18-64, and 65+. | Yes | Improvement Over Self | 2019 | 2A. Bi-Directional Integration 2C. Transitional Care 3A. Opioid Use 3D. Chronic Disease Prevention |
| Patients on high-dose chronic opioid therapy by varying thresholds | Measure specification in development. Among Medicaid enrollees, the percentage of chronic opioid therapy patients receiving doses: >50 mg. MED in a quarter, doses >90 mg. MED in a quarter. | No | Improvement Over Self | 2019 | 3A. Opioid Use |
| Patients with concurrent sedatives prescriptions | Measure specification in development. Among Medicaid enrollees receiving chronic opioid therapy, the percentage that had more than 45 days of Sedative Hypnotics/ Benzodiazepines/ | No | Improvement Over Self | 2019 | 3A. Opioid Use |

| Measure Name | Measure Description | Statewide DSRIP Accountability Measure? | Method for Assessment of ACH Performance | Initial Assessment Year | Associated Project Areas |
|--|--|---|--|-------------------------|---|
| | carisoprodol/ barbiturates dispensed in the quarter. | | | | |
| Percent Homeless (Narrow Definition) | Percent of Medicaid enrollees who were homeless in at least one month in the measurement year. Excludes “homeless with housing” ACES living arrangement code | No | Improvement Over Self | 2019 | 2C. Transitional Care |
| Plan All-Cause Readmission Rate (30 Days) | The proportion of acute inpatient stays during the measurement year that were followed by an unplanned acute readmission within 30 days among Medicaid enrollees ages 18-64 years old. | Yes | Improvement Over Self | 2019 | 2A. Bi-Directional Integration 2C. Transitional Care |
| Statin Therapy for Patients with Cardiovascular Disease (Prescribed) | Percentage of male Medicaid enrollees 21 to 75 years of age and female Medicaid enrollees 40 to 75 years of age during the measurement year who were identified as having clinical ASCVD who were dispensed at least one high- or moderate-intensity statin medication. | No | Improvement Over Self | 2020 | 3D. Chronic Disease Prevention |
| Substance Use Disorder Treatment Penetration | The percentage of Medicaid enrollees with a substance use disorder treatment need who received substance use disorder treatment in the measurement year Separate reporting by age groups: <ul style="list-style-type: none"> • 12-17 years; • 18-64 years; | Yes | Improvement Over Self | 2019 | 2A. Bi-Directional Integration |

| Measure Name | Measure Description | Statewide DSRIP Accountability Measure? | Method for Assessment of ACH Performance | Initial Assessment Year | Associated Project Areas |
|---|--|---|--|-------------------------|--------------------------|
| | <ul style="list-style-type: none"> 65+ years. | | | | |
| Substance Use Disorder Treatment Penetration (Opioid) | Measure specification in development. Percent of Medicaid enrollees with a diagnosis of opioid use disorder who have a substance use service need who received at least one qualifying service during the measurement year. Reported separately for adults and for children. | No | Improvement Over Self | 2020 | 3A. Opioid Use |